



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

Block Level Experimental Agromet Advisory Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



CHALAKUDY BLOCK, THRISSUR DISTRICT

AAS 26/2018(52), Friday

29.06.2018

SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:

The total rainfall received 81.5 mm. the maximum temperature ranges from 28.8°C to 31.4°C, minimum temperature ranges from 21.9°C to 23.5°C. Morning relative humidity ranges from 95% to 98% and afternoon relative humidity ranges from 70% to 98%. Wind speed ranges from 0.6 to 1.9 km/h. Bright sunshine hours range from 0.0 to 4.5 hours. Evaporation ranges from 1.4 to 3.1 mm.

CURRENT SYNOPTIC SITUATION: A cyclonic circulation lies over east central Arabian Sea off south Maharashtra coast at 7.6 km above mean sea level.

Heavy Rainfall warning: Day 1 & 2: Heavy rainfall is most likely to occur at one or two places in Kerala.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	30.06.2018	01.07.2018	02.07.2018	03.07.2018	04.07.2018
Rainfall (mm)	10	6	2	8	11
Maximum temperature (°C)	28	29	29	29	29
Minimum temperature (°C)	22	23	23	23	23
Total cloud cover (octa)	8	6	5	8	7
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	015	015	015	015	015
Wind direction (deg)	250	250	250	250	250

Below Normal rainfall is expected in Kerala for the coming week.

Moderate rainfall is expected in Chalakudy block for next five days.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Crop stage	Pest/ Disease	Advisories
Banana	Bunch emergence stage	Erwinia Rot	Proper drainage should be ensured. As a precaution, apply lime in the banana basin or bleaching powder in the irrigation channels. To improve the disease resistance, drench the basin with Pseudomonas @ 20g per litre of water. Affected plants can be protected by drenching the basin with 5 g bleaching powder or 3 g copper oxychloride or 2g copper hydroxide per one litre of water. Severely affected plants should be cut and removed to prevent the spread of disease.
Vegetables (cowpea)	-	Fusarium wilt	Due to continuous rainfall and water logging, there is a chance of fusarium wilt in cowpea. Drain the field properly and drench the soil rhizosphere with 2 g of bavistin or 2 ml of contaf per one litre of water alternatively within 15 days interval to control the disease. Severely affected plants should be removed and destroyed from the field and apply a pinch of bleaching powder in the pit.
Animal Husbandry	Avoid grazing of the animals in waterlogged areas. Farmers should be aware of leptospirosis. Contact nearest veterinary hospital for HS and FMD vaccination. (CAADECCS, KVASU, Mannuthy)		