



## Agromet Advisory Service bulletin 02/16, Tuesday dt 02-02-16

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

## **ERNAKULAM**

## **CURRENT SYNOPTIC SITUATION:**

A trough of low pressure at mean sea level lies over South West Bay of Bengal and adjoining Sri Lanka coast

## NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 02.02.2016 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS		ENSEMBLE FCST			
	DAY-1 03/02	DAY-2 04/02	DAY-3 05/02	DAY-4 06/02	DAY-5 07/02
DISTRICT : ERNAKULAM					
Rainfall (mm)	0	0	0	0	2
Max Temperature ( deg C)	32	32	32	32	32
Min Temperature ( deg C)	23	23	23	23	23
Total cloud cover (octa)	5	5	5	6	5
Max Relative Humidity (%)	80	80	80	80	80
Min Relative Humidity (%)	55	55	55	55	55
Wind speed (kmph)	800	008	800	008	800
Wind direction (deg)	270	270	270	270	270

Agro Meteorological Advisories

Crop	Pest/Disease	Advisories	
Paddy		Punja is in seedling stage.	
Coconut	Root wilt	There is a chance of Root wilt in coconut garden. Apply 50 kg cattle manure or green manure and 1kg of lime/palm/year. Growing green manure crops like sun hemp, sesbania, cowpea in the coconut basin and their incorporation in situ is beneficial as the practice reduce the intensity of the root wilt and increase the nut yield. Rogue out palms that are affected severely by root wilt and yield less than 10 nuts/palm/year and those. Replant with disease tolerant high yielding hybrids.	
Banana	Cigar end disease  Due to high temperature and low humidity, there is a character of Cigar end disease in banana bunches. Spray 3 g of Dithan @ 1 liter of water. The banana plantation may be given so that the plantation can with stand high wind speed.		
Vegetables	Sukking pests	Due to high temperature experienced, there is a chance of attack of white fly, hopper and aphid in vegetables. Apply 20 g Verticillium per one liter of water. 5 ml sunflower oil or castor oil along with 10 g of jaggery can be added to the spray solution to make the spray effective. Otherwise apply tobacco decoction.	