





## Agromet Advisory Bulletin for Ernakulam District (Valid from 14.5.2022 to 18.5.2022)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 19/2022, (38) Friday dated 13.5.2022

## A. Medium range weather forecast of IMD for next five days

RAINFALL FORECAST FOR NEXT FIVE DAYS							
Weather parameters		14.5.2022	15.5.2022	16.5.2022	17.5.2022	18.5.2022	
Rainfall (mm)		15	20	20	7	5	
Maximum temperature (°C)		30	31	32	33	33	
Minimum temperature (°C)		24	25	26	27	27	
Maximum Relative Humidity (%)		90	90	90	90	90	
Minimum Relative Humidity (%)		85	85	85	85	85	
Wind speed (kmph)		8	10	8	4	4	
Wind direction (deg)		270	270	270	270	270	
Total cloud cover (octa)		7	8	8	8	7	
2022 May 14	2022 May 15	2022 May 16 202		2 May 17	2022	May 18	
Light rain	Moderate rain	Moderate rain Li		ght rain	Ligh	t rain	

**Weather summary / Alert:** Moderate rainfall is expected in the district and heavy rainfall is expected in isolated places. Above normal rainfall is expected in Kerala.

## B. Agrometeorological Advisories for crop protection & management

Ensure drainage facilities in low land. Proper mulching can be followed to conserve soil moisture.

C. General and Agrometeorological Advisories for Ernakulam District						
Crop	Pest/ Dise	Crop/ A	Crop/ Animal Advisory(RRS Vyttila)			
Paddy (Pokkali)	Management	nter bunds. During low tide, v dry. Close the shutters of	operations in pokkali fields. Strengthen the water has to be drained out of the field and let the sluice after that. Take mounds of 1 mer drying the field and leave it as such for the			
Coconut	Leaf rot	Improve general condition of the palm by manuring and management. Clean the crown and remove rotten portions. Spray crowns and leaves with 1 % Bordeaux mixture, Mancozeb 75 WP 3g per litre water, Copper Oxy Chloride 50 WP 3g per litre.				
Banana	Management	Apply last dose of fertilizer along with earthing up before bunch emergence. Propping of plants also recommended.				
Mango	Fruit flies	Destroy all fallen fruits at weekly intervals. Install six methyl eugenol plywood traps (pheromone traps) per acre. Plough the soil at the tree basin at frequent intervals.				
Fisheries	It is the right time to prepare ponds for fish farming. Drain the pond and apply lime. Apply tea seed cake or mahua oil cake in ponds where draining is difficult to eradicate weed fishes.					