

Agromet Advisory Bulletin for the District, Kasaragod

(Valid from 29.03.2024 to 02.04.2024)



(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode& India Meteorological Department)

A. Weather Summary of preceding four days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
0.0	34.0 – 34.5	25.0 – 25.4	63 – 81	04 - 05

B. Weather forecast for next five days

Parameters	29-03-2024	30-03-2024	31-03-2024	01-04-2024	02-04-2024
Rainfall, mm	0	0	0	0	0
Max. Temp, °C	35	35	35	35	35
Min. Temp,°C	26	26	26	26	26
Max. Relative Humidity, %	85	85	85	85	85
Min. Relative Humidity, %	59	59	59	59	59
Wind speed,km/h	4	4	4	2	2
Wind direction, degrees	290	290	290	250	250
Total cloud cover, octa	4	2	2	3	2

C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories		
General	No rain **				
conditions	No rainfall. Temperatures will be higher during the day. Atmospheric humidity will be normal.				
General Recommen dations	Mulch the crop basins. Irrigate the crop when the water is available in the evening or early morning. Adopt drip irrigation method for maximum water use efficiency. 1. Arrange for irrigation facilities from available water resources. 2. Remove weeds from the soil to reduce transpiration losses. Powder the soil to dust by breaking the clods. This will act as good soil mulch to prevent evaporation loss of water. 3. Well drained areas where lifesaving irrigation possible ragi and millets cause cultivated. 4. Take care of controlling of sucking pests; control/minimize the insect and pest incidence with IPM.				
	5. Repair and rejuvenate local water bodies before the rainy season.				

Various crops	Various stages	Sucking pests The climate is favourable for the spread of sucking pests like mealy bug, jasids, aphids, mites, bugs etc. If not controlled properly they will act as vectors and may spread virus diseases.	To control the pests apply neem oil emulsion (5 ml. neem oil mixed in one litre of luke warm soap water solution) Or Apply malathion 50 EC @ 2 ml + neem oil 4ml per litre of water
Coconut	All stages	Drought Management	 Cut two green leaves from the bottom layer, to reduce the water loss from the tree. Apply compost/dried leaves in the basins to increase water holding capacity. Adopt drip irrigation. This will minimize the irrigation water loss. Take care of controlling of sucking pests; control/minimize the insect and pest incidence with IPM.
Coconut	All stages	Stem bleeding	Reduce the nut load by harvesting all the matured and about to matured nuts Chisel out the affected parts and apply Hexaconazole (Contaf®) 5ml/litre solution in the wound. After getting the wound dried, immediately smear the spot with Rubbercoat®. Drench the palm basin (2m radius) with Hexaconazole (Contaf®), 50ml dissolved in 25 litre of water per palm at monthly intervals, for 5months or till the disease fully disappears. Spread lime thickly in the basins and incorporate with the soil.

Black pepper	Different stages	Wilting due to drought	Protect the plants from direct sunlight by smothering the basal portions of the vines with dried banana leaves, plated coconut leaves etc Give protective irrigations one in every two weeks.
Cucurbitaceo us vegetables	All stages	Downy mildew	Downy mildew: As a prophylactic measure apply 'Mancozeb'(@ 2g/l of water). If disease appeared, spray Akomin® (@3ml/L) on both surfaces of the leaves, thrice at 15 days intervals.
Poultry and pet birds	Different stages	Summer stress	To combat heat stress, the poultry sheds should be protected from direct sunlight, roofing can be painted white to reflect heat, fans can be fitted, cool water can be sprayed, plenty of clean water can be provided with ice, glucose and 0.1 % sodium bicarbonate, feed offered during the cooler parts of the day can be supplemented with 20% extra vitamins, phosphorous and vitamin C.
Animal Husbandry	All stages	Summer Stress	The rise in temperature will affect the thermoregulatory mechanism of dairy cattle. This will cause increase in body temperature, rapid shallow breathing, increased heart rate, profuse salivation, and reduced feed intake. This in turn results in severe production loss and reduced breeding efficiency in dairy cattle.
			Provide pure drinking water to the dairy cattle (45 to 60 litres of water), Allow grazing only during the cooler parts of the day. Provide shading. Shelter them in thatched roofings of minimum 9 ft. height with ample ventilation. Providing fans, misting and fogging assembly in cattle sheds will help them to regulate body temperature. Also ensure minerals fortified feeds.

Cow Milking Summer Mastitis

Summer Mastitis is a bacterial disease. The disease is common on cows which are sheltered in unhygienic conditions. The disease is transmitted by insects like flies and mosquitoes.

Symptoms:- The symptoms are change in composition of milk and turning the milk into an yellow watery liquid, severe fever, lack of appetite, abortions etc. If not treated the cow may lose its milk yielding capacity permanently. Also the cow may lose mobility and gradually death may occur.

Control:- Ensure hygiene cattle shed and surroundings to control the pathogen and the vectors. Boost the immunity of cattle by supplementing feeds with vitamins and minerals containing health tonics.

If disease appears, immediately give medication under the supervision of a Veterinary doctor.

** Warning colour codes of rainfall (for disaster management)

Warning (Take actions)

Alert (Be prepared)

Watch (Be updated)

No warning (No actions)

Sd/-Nodal Officer, GKMS Project, RARS Pilicode