



## Agromet Advisory Bulletin for PALAKKAD District

(Valid from 22.01.2025 to 26.01.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara  
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 03/2025, (6) Tuesday

Dated 21.01.2025

### A. Medium range weather forecast of IMD for next five days

Weather parameters	22.01.2025	23.01.2025	24.01.2025	25.01.2025	26.01.2025
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	33	33	33	33	33
Minimum temperature (°C)	24	24	24	24	24
Maximum Relative Humidity (%)	82	82	82	82	82
Minimum Relative Humidity (%)	46	46	46	46	46
Wind speed (kmph)	4	3	2	2	4
Wind direction (deg)	70	20	50	70	70
Total cloud cover (octa)	2	2	3	3	2

**Weather summary / Alert:** Since dry weather condition is prevails, irrigation of crops can be continued as per requirement. Normal to above normal rainfall is expected in Kerala.

### B. Agrometeorological Advisories for crop protection & management

#### General and Agrometeorological Advisories for Thrissur District

**General Advisory:** - Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Rice (Mundakan) Dough stage	Bacterial Leaf Blight	To control and prevent Bacterial leaf blight in paddy, dissolve 20 g of cow dung in one litre of water and spray the supernatant liquid by mixing with 20g of pseudomonas. If the attack is severe spray 6 g of Streptocycline in 20 litre of water.
Banana	Management	Since the wind speed may increase, give strong propping to banana plants.
Bhindi	Jassid's	Due to increase in atmospheric temperature, the attack of Jassid's in Bhindi was increased. They can be controlled by applying 2% neem oil emulsion once in a week or by spraying 2 g thiamethoxam per ten litre of water.
Cowpea	Pod borer	There is a chance of pod borer attack in cowpea. It can be controlled by spraying 50 grams Neem kernel extract per one liter of water. Or spray 20 grams of Beauveria in one liter of water.

**Animal Husbandry:** Contact nearby veterinary hospitals to vaccinate the animals against infectious diseases. Care should be taken to avoid fungal growth in the cattle feed. (CAADECCS, KVASU, Mannuthy).