Agromet Advisory Service bulletin 19/16 Tuesday dt.03-05-2016

(Issued jointly by IAAS Vellanikkara)


Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

THRISSUR

Significant weather for the preceding week at Vellanikkara



Rainfall weekly total	0.0 mm
Maximum temperature	36.3-36.9 °C
Minimum temperature	25.5-27.2° C
Max. RH	78-89%
Min. RH	50-58%
Wind speed	1.8-2.7 km/hr
Sun shine hour	6.8-10.0 hr
Evaporation	5.5-4.6 mm



CURRENT SYNOPTIC SITUATION:

An upper air trough runs from Telangana to Comorin area, extends upto 0.9km above sea level.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 03.05.2016 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS		ENSEM	ENSEMBLE FCST		
	DAY-1 04/05	DAY-2 05/05	DAY-3 06/05	DAY-4 07/05	DAY-5 08/05
DISTRICT : THRISSUR					
Rainfall (mm)	0	3	4	3	2
Max Temperature (deg C)	36	36	36	35	35
Min Temperature (deg C)	26	26	25	25	25
Total cloud cover (octa)	4	4	4	6	6
Max Relative Humidity (%)	80	80	80	85	85
Min Relative Humidity (%)	60	60	60	65	65
Wind speed (kmph)	006	006	006	006	006
Wind direction (deg)	250	250	250	250	230

Agro Meteorological Advisories

Crop	Stage of the crop /	Advisories
	Pest/Disease	
Rice (Harvesting stage)	Fungal diseases (Blast & Sheath blight)	Apply Pseudomonas culture (2.5 kg/acre). Apply organic fertilizers. In severely affected areas, spray Flusilazole or Propiconasole @ 1 ml per litre of water.
Coconut	Rhinoceros beetle	Apply 250 gm neem cake mixed with equal volume of sand in the innermost 2-3 leaf axils.
Banana	Yellowing (Erwinia Rot)	There is a chance of yellowing in banana. Examine the sucker and make sure that it is Erwinia rot, it can be controlled by mixing 25gms of Kocide and 6gms of Streptocycline in 25 ml of water and apply 5 litre of solution to each banana basin.

Vegetables Mosaic	There is a chance of mosaic disease in vegetables. Control the vectors by spraying dimethoate 0.05	
vegetables	Wosaic	percent. Uprooting and destruction of affected plants
		and collateral hosts should be done.