



Agromet Advisory Bulletin for Thrissur District

(Valid from 09.07.2025 to 13.07.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 27/2025, (54) Tuesday

Dated 08.07.2025

| Weather parameters | 09.07.2025 | 10.07.2025 | 11.07.2025 | 12.07.2025 | 13.07.2025 |
|-------------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 12 | 8 | 10 | 10 | 3 |
| Maximum temperature (°C) | 30 | 30 | 30 | 30 | 30 |
| Minimum temperature (°C) | 23 | 23 | 23 | 23 | 23 |
| Maximum Relative Humidity (%) | 90 | 90 | 90 | 90 | 90 |
| Minimum Relative Humidity (%) | 85 | 85 | 85 | 85 | 85 |
| Wind speed (kmph) | 3 | 3 | 3 | 3 | 3 |
| Wind direction (deg) | 340 | 340 | 340 | 340 | 340 |
| Total cloud cover (octa) | 8 | 7 | 8 | 8 | 8 |

Weather summary / Alert: - As per the forecast of IMD, there is a chance of light rainfall in the district and moderate rainfall in isolated places of the district. Strong winds speed reaching likely to prevail occasionally over Kerala from 08th to 10th July 2025. Above normal rainfall is expected in Kerala.

General and Agrometeorological Advisories for Thrissur District

General Advisory: - Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. Pesticides should not be applied during rainy period. Provide supports for the bananas. Make the vegetable pandals stronger. Avoid standing in open areas to protect farmers and livestock from lightning and thunder.

| Crop | Pest/ Disease, Crop Management | Agromet Advisory |
|---------------------------|--------------------------------|---|
| Rice (virippu) | Thrips | To control thrips in rice, spray 2 gram of Thiomethoxam in 10 litre of water. Pesticides should not be applied during rainy period. |
| Pepper | Quick wilt | During rainy season, as a prophylactic measure to control quick wilt of pepper, add 2 kg Trichoderma to a mixture of 90 kg FYM and 10 kg neem cake and kept for two weeks with sufficient moisture. Add 2.5 kg of this mixture to the base of each matured pepper vine. If the attack is severe, drench the basin of each plant with three gram Copper Oxy Chloride (COC) per one litre of water. |
| Cucurbitaceous Vegetables | Downy mildew | Proper drainage facilities have to be ensured. Pesticides should not be applied during rainy period. During rainy season, there is a chance of downy mildew attack in cucurbitaceous vegetables. Spray 2.5gm Mancozeb per one litre of water under the leaves. |

Animal Husbandry: Sundry the cattle feed frequently to avoid fungal growth. Give foot and mouth disease (FMD) vaccination to the animal. Don't allow the animals go to water logged areas in order to avoid anthrax and foot lesions. (CAADECCS, KVASU, Mannuthy)